

ORIGINAL ARTICLE

Lecanemab in Early Alzheimer's Disease

Christopher H. van Dyck, M.D., Chad J. Swanson, Ph.D., Paul Aisen, M.D., Randall J. Bateman, M.D., Christopher Chen, B.M., B.Ch., Michelle Gee, Ph.D., Michio Kanekiyo, M.S., David Li, Ph.D., Larisa Reyderman, Ph.D., Sharon Cohen, M.D., Lutz Froelich, M.D., Ph.D., Sadao Katayama, M.D., Marwan Sabbagh, M.D., Bruno Vellas, M.D., David Watson, Psy.D., Shobha Dhadda, Ph.D., Michael Irizarry, M.D., Lynn D. Kramer, M.D., and Takeshi Iwatsubo, M.D.et al.

November 29, 2022

DOI: 10.1056/NEJMoa2212948

Article

Figures/Media

Metrics

27 References

Abstract

BACKGROUND

The accumulation of soluble and insoluble aggregated amyloid-beta ($A\beta$) may initiate or potentiate pathologic processes in Alzheimer's disease. Lecanemab, a humanized IgG1 monoclonal antibody that binds with high affinity to $A\beta$ soluble protofibrils, is being tested in persons with early Alzheimer's disease.

METHODS

We conducted an 18-month, multicenter, double-blind, phase 3 trial involving persons 50 to 90 years of age with early Alzheimer's disease (mild cognitive impairment or mild dementia due to Alzheimer's disease) with evidence of amyloid on positron-emission tomography (PET) or by cerebrospinal fluid testing. Participants were randomly assigned in a 1:1 ratio to receive intravenous lecanemab (10 mg per kilogram of body weight every 2 weeks) or placebo. The primary end point was the change from baseline at 18 months in the score on the Clinical Dementia Rating–Sum of Boxes (CDR-SB; range, 0 to 18, with higher scores indicating greater impairment). Key secondary end points were the change in amyloid burden on PET, the score on the 14-item cognitive subscale of the Alzheimer's Disease Assessment Scale

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



on the Alzheimer's Disease Cooperative Study–Activities of Daily Living Scale for Mild Cognitive Impairment (ADCS-MCI-ADL; range, 0 to 53; lower scores indicate greater impairment).

RESULTS

A total of 1795 participants were enrolled, with 898 assigned to receive lecanemab and 897 to receive placebo. The mean CDR-SB score at baseline was approximately 3.2 in both groups. The adjusted least-squares mean change from baseline at 18 months was 1.21 with lecanemab and 1.66 with placebo (difference, -0.45 ; 95% confidence interval [CI], -0.67 to -0.23 ; $P < 0.001$). In a substudy involving 698 participants, there were greater reductions in brain amyloid burden with lecanemab than with placebo (difference, -59.1 centiloids; 95% CI, -62.6 to -55.6). Other mean differences between the two groups in the change from baseline favoring lecanemab were as follows: for the ADAS-cog14 score, -1.44 (95% CI, -2.27 to -0.61 ; $P < 0.001$); for the ADCOMS, -0.050 (95% CI, -0.074 to -0.027 ; $P < 0.001$); and for the ADCS-MCI-ADL score, 2.0 (95% CI, 1.2 to 2.8 ; $P < 0.001$). Lecanemab resulted in infusion-related reactions in 26.4% of the participants and amyloid-related imaging abnormalities with edema or effusions in 12.6%.

CONCLUSIONS

Lecanemab reduced markers of amyloid in early Alzheimer's disease and resulted in moderately less decline on measures of cognition and function than placebo at 18 months but was associated with adverse events. Longer trials are warranted to determine the efficacy and safety of lecanemab in early Alzheimer's disease. (Funded by Eisai and Biogen; Clarity AD ClinicalTrials.gov number, [NCT03887455](https://clinicaltrials.gov/ct2/show/study/NCT03887455).)

Introduction

CURRENT THERAPEUTIC AGENTS FOR ALZHEIMER'S DISEASE–RELATED DEMENTIA temporarily improve symptoms but do not alter the underlying disease course.^{1,2} Some evidence suggests that amyloid removal slows the progression of disease.³ One anti-amyloid antibody (aducanumab) has received accelerated approval from the Food and Drug Administration.

Lecanemab is a humanized monoclonal antibody that binds with high affinity to soluble amyloid-beta ($A\beta$) protofibrils, which have been shown to be more toxic to neurons than monomers or insoluble fibrils.⁴⁻¹⁴ A phase 2b, dose-finding trial involving 854 participants with early Alzheimer's disease did not show a significant difference between lecanemab and placebo in a Bayesian analysis of 12-month change in a composite score (primary end point). However, analyses at 18 months showed dose- and time-dependent clearance of amyloid with lecanemab, and the drug was associated with less clinical decline on some measures than placebo. In that trial, intravenous administration of 10 mg of lecanemab per kilogram of body weight every 2 weeks was identified as an appropriate dose, with a 9.9% incidence ($< 3\%$ symptomatic) of amyloid-related imaging abnormalities (ARIA) with edema or effusions (ARIA-E).¹⁵ We conducted a phase 3 trial (Clarity AD) to determine the safety and efficacy of lecanemab in participants with early Alzheimer's disease.

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



Methods



TRIAL DESIGN AND OVERSIGHT

Clarity AD was an 18-month, multicenter, double-blind, placebo-controlled, parallel-group trial involving persons with early Alzheimer's disease. Eligible participants were randomly assigned in a 1:1 ratio to receive intravenous lecanemab (10 mg per kilogram every 2 weeks) or placebo. The randomization was stratified according to clinical subgroup (mild cognitive impairment due to Alzheimer's disease or mild Alzheimer's disease–related dementia on the basis of the criteria noted below), the presence or absence of concomitant approved medication for symptoms of Alzheimer's disease at baseline (e.g., acetylcholinesterase inhibitors, memantine, or both), apolipoprotein E (ApoE) ϵ 4 carriers or noncarriers, and geographic region. During the trial, participants underwent serial blood testing for plasma biomarkers and could participate in three optional substudies that evaluated longitudinal changes in brain amyloid burden as measured by positron-emission tomography (PET), brain tau pathologic features as measured by PET, and cerebrospinal fluid (CSF) biomarkers of Alzheimer's disease.

The trial was conducted in accordance with International Council for Harmonisation guidelines and the ethical principles of the Declaration of Helsinki. The trial was approved by the institutional review board or independent ethics committee at each center, and all the participants provided written informed consent. The sponsor Eisai designed the trial and analyzed the data in collaboration with the academic authors, provided lecanemab and placebo, provided funding for medical writing, and aided in drafting the manuscript. The sponsor could not delay or interdict publication. The first, second, and sixteenth authors wrote the first draft of the manuscript, with professional medical writing assistance funded by Eisai, and all the authors contributed to subsequent drafts. Confidentiality agreements were in place between the sponsor and the authors and site investigators. Biogen provided partial funding for the trial.

An independent data and safety monitoring board consisting of experts in Alzheimer's disease and statistics reviewed unblinded safety data during the trial. An independent medical monitoring team, whose members were unaware of the trial-group assignments, reviewed ARIA, infusion-related reactions, and hypersensitivity reactions. Clinical assessment raters were unaware of the safety assessments and the trial-group assignments. All the authors vouch for the completeness and accuracy of the data, the fidelity of the trial to the [protocol](#) (available with the full text of this article at NEJM.org), and the full reporting of adverse events.

ELIGIBILITY CRITERIA

The trial included participants 50 to 90 years of age, with either mild cognitive impairment due to Alzheimer's disease or mild Alzheimer's disease–related dementia on the basis of National Institute on Aging–Alzheimer's Association criteria.^{16,17} Amyloid positivity was determined by PET or CSF measurement of A β ₁₋₄₂. All the participants had objective impairment in episodic memory as indicated by at least 1 standard deviation below the age-adjusted mean in the Wechsler Memory Scale IV–Logical

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



END POINTS

The primary efficacy end point was the change in the score on the Clinical Dementia Rating (CDR)–Sum of Boxes (CDR-SB)¹⁸ from baseline at 18 months. The CDR-SB score is a validated outcome measure used in clinical trials of Alzheimer's disease that is obtained by interviewing patients and their care partners and captures cognition and function. It assesses six domains that patients and caregivers identify as important (Memory, Orientation, Judgment and Problem Solving, Community Affairs, Home and Hobbies, and Personal Care). Scores for each domain range from 0 to 3, with higher scores indicating greater impairment. Total scores range from 0 to 18, with a score of 0.5 to 6 indicating early Alzheimer's disease.

Key secondary end points were the change from baseline at 18 months in the following: amyloid burden on PET as measured in centiloids (with either florbetaben, florbetapir, or flutemetamol tracers) in a substudy, the score on the 14-item cognitive subscale of the Alzheimer's Disease Assessment Scale (ADAS-cog14; range, 0 to 90, with higher scores indicating greater impairment),¹⁹ the Alzheimer's Disease Composite Score (ADCOMS; range, 0 to 1.97, with higher scores indicating greater impairment),²⁰ and the Alzheimer's Disease Cooperative Study–Activities of Daily Living Scale for Mild Cognitive Impairment (ADCS-MCI-ADL; range, 0 to 53, with lower scores indicating greater impairment).²¹ Biomarker assessments included CSF biomarkers ($A\beta_{1-40}$, $A\beta_{1-42}$, total tau, phosphorylated tau 181 [p-tau181], neurogranin, and neurofilament light chain [NfL]) and plasma biomarkers ($A\beta_{42/40}$ ratio, p-tau181, glial fibrillary acidic protein [GFAP], and NfL). Tau PET and volumetric magnetic resonance imaging (MRI) results have not been fully analyzed.

A prespecified exploratory and multiplicity-unadjusted analysis examined the time to worsening of the global CDR score (range, 0 to 3, with higher scores indicating greater impairment). This end point was defined as the time to the first increase of at least 0.5 points in the global CDR score on two consecutive visits.

STATISTICAL ANALYSIS

The sample size for this trial was estimated on the basis of comparison of lecanemab and placebo with respect to the primary efficacy end point, the change from baseline at 18 months in the CDR-SB score. On the basis of data from the phase 2b trial of lecanemab,¹⁵ the estimated standard deviation of the change from baseline at 18 months in the CDR-SB score with placebo was 2.031 points, and the estimated treatment difference between lecanemab and placebo in all the participants was 0.373 points. This estimation corresponds to 25% less decline in cognitive function with lecanemab than with placebo and is consistent with a clinically meaningful difference on the basis of the Alzheimer's disease literature, statistical principles, and agreements with regulatory authorities.^{15,22-24} Therefore, under the assumption of an estimated 20% dropout rate at 18 months in this trial, a total sample size of 1566 participants, including 783 participants receiving lecanemab and 783 participants receiving placebo, would provide the trial with 90% power to detect the treatment difference with the use of a two-sample t-test at a two-sided alpha level of 0.05. The sample size was increased by 200 to account for participants who missed three or more consecutive doses during the initial 6-month peak period of

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



Efficacy analyses were performed in the modified intention-to-treat population, which was defined as the group of randomly assigned participants who received at least one dose of lecanemab or placebo and who had a baseline assessment and at least one postdose primary efficacy (CDR-SB) measurement. Sensitivity analyses across efficacy end points to assess the robustness of the primary analysis to missing data included rank analysis of covariance with imputation of missing values. Additional sensitivity analyses were performed to evaluate potential effects of functional unblinding due to ARIA and effects of missed doses due to Covid-19–related absences (see the [Supplementary Appendix](#), available at NEJM.org). Safety was evaluated in the safety population, which was defined as the group of participants who received at least one dose of lecanemab or placebo. Safety evaluations included monitoring of adverse events, vital signs, physical examinations, clinical laboratory variables, and 12-lead electrocardiograms. Occurrences of ARIA were monitored throughout the trial by central reading of MRI performed at weeks 9, 13, 27, 53, and 79 as well as at the 3-month follow-up visit (week 91) for safety monitoring. In addition, the populations for the substudies of amyloid burden on PET, tau pathologic features on PET, and CSF biomarkers of Alzheimer's disease were the groups of participants who received at least one dose of lecanemab or placebo and who underwent a baseline PET or CSF evaluation and at least one postdose evaluation.

The primary analysis was performed without imputation of missing values. The primary analysis of the change from baseline at 18 months in the CDR-SB score was performed to compare lecanemab and placebo with the use of a mixed model for repeated measures that included the baseline CDR-SB score as a covariate, with trial group, visit, stratification variables (i.e., clinical subgroup, use of medication for symptoms of Alzheimer's disease at baseline [yes or no], ApoE ϵ 4 carrier status [carriers or noncarriers], and geographic region [North America, Europe, and Asia–Pacific]), baseline CDR-SB score–by–visit interaction, and trial group–by–visit interaction as fixed effects. If the between-group difference in primary end-point results was significant, then key secondary end points were to be tested hierarchically in the following order: change from baseline at 18 months in amyloid burden on PET as measured in centiloids in the subgroup tested and change from baseline at 18 months in the ADAS-cog14 score, change from baseline at 18 months in the ADCOMS, and change from baseline at 18 months in the ADCS-MCI-ADL score, all in the modified intention-to-treat population. Each test was performed at an alpha level of 0.05 (two-sided) and was to be performed only if the preceding test was significant at a two-sided level of 0.05. Additional details on the design and analysis methods are provided in the [Supplementary Appendix](#) and protocol.

Results




PARTICIPANTS

Figure 1.

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.





Screening, Randomization, and Follow-up.

Table 1.

Characteristic	Lecanemab (N=859)	Placebo (N=875)
Age — yr	71.4±7.9	71.0±7.8
Sex — no. (%)		
Female	443 (51.6)	464 (53.0)
Male	416 (48.4)	411 (47.0)
Race — no. (%) [†]		
White	655 (76.3)	677 (77.4)
Black	20 (2.3)	24 (2.7)
Asian	147 (17.1)	148 (16.9)
Other or missing	37 (4.3)	26 (3.0)
Hispanic ethnic group — no. (%) [‡]	107 (12.5)	108 (12.3)
Time since diagnosis — yr	1.4±1.51	1.3±1.54
Time since onset of symptoms — yr	4.13±2.35	4.15±2.53
Global CDR score — no. (%) [‡]		
0-5	694 (80.8)	706 (80.7)
1	165 (19.2)	169 (19.3)
Clinical subgroup — no. (%)		
Mild dementia due to Alzheimer's disease	331 (38.5)	331 (37.8)
Mild cognitive impairment due to Alzheimer's disease	528 (61.5)	544 (62.2)
ApoE ε4 status — no. (%)		
Noncarrier	267 (31.1)	275 (31.4)
Carrier	592 (68.9)	600 (68.6)
Heterozygotes	456 (53.1)	468 (53.5)
Homozygotes	136 (15.8)	132 (15.1)
Current use of medication for symptoms of Alzheimer's disease — no. (%)	447 (52.0)	468 (53.5)
CDR-SB score [‡]		
Mean	3.17±1.34	3.22±1.34
Range	0.5 to 8.0	0.5 to 8.5
Amyloid burden on PET — centiloids [¶]		
Mean	77.92±44.84	75.03±41.82
Range	-16.6 to 213.2	-17.0 to 179.6
ADAS-cog14 score [‡]		
Mean	24.45±7.08	24.37±7.56
Range	4.7 to 47.7	5.0 to 60.7
ADCCMS ^{**}		
Mean	0.398±0.147	0.400±0.147
Range	0.08 to 0.94	0.07 to 0.91
ADCS-MCI-ADL score [‡]		
Mean	41.2±6.6	40.9±6.9
Range	13 to 53	12 to 53
MMSE score [‡]		
Mean	25.5±2.2	25.6±2.2
Range	22 to 30	22 to 30

^a Plus-minus values are means ±SD. ApoE denotes apolipoprotein E.
[†] Race and ethnic group were determined by the participants.
[‡] Global Clinical Dementia Rating (CDR) scores range from 0 to 3, with higher scores indicating greater impairment. A score of 0.5 is considered to be the threshold for Alzheimer's disease and was required for trial enrollment.
[§] Scores on the CDR-Sum of Boxes (CDR-SB) range from 0 to 18, with higher scores indicating greater impairment.
[¶] Values for amyloid burden on positron-emission tomography (PET) were for the PET substudy population.
^{**} Scores on the 14-item cognitive subscale of the Alzheimer's Disease Assessment Scale (ADAS-cog14) range from 0 to 90, with higher scores indicating greater impairment.
^{††} Values for the Alzheimer's Disease Cooperative Study-Activities of Daily Living Scale for Mild Cognitive Impairment (ADCS-MCI-ADL) range from 0 to 53, with lower scores indicating greater impairment.
^{‡‡} Scores on the Mini-Mental State Examination (MMSE) range from 0 to 30, with lower scores indicating greater impairment.

Characteristics of the Participants at Baseline (Modified Intention-to-Treat Population).

A total of 5967 persons were screened and 1795 underwent randomization; 898 were assigned to receive lecanemab and 897 to receive placebo at 235 sites in North America, Europe, and Asia from March 2019 through March 2021. Of these participants, 729 (81.2%) in the lecanemab group and 757 (84.4%) in the placebo group completed the trial and had data available on the primary end point (Figure 1). The modified intention-to-treat population included 1734 participants (859 in the lecanemab group and 875 in the placebo group), and the safety population included all 1795 randomly assigned participants. Enrollment in three longitudinal substudies included 698 participants in the substudy of amyloid burden on PET, 757 in the study of tau pathologic features on PET, and 781 in the substudy of CSF


This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



group of participants (20% non-White), including in the United States, where 6.1% and 28.1% of the 3638 screened participants and 4.5% and 22.5% of randomly assigned participants were Black and Hispanic, respectively. The characteristics of the participants at baseline were generally similar in the two trial groups (Table 1). These characteristics were similar to what has been observed in population studies involving persons with early Alzheimer's disease, although there was an underrepresentation of Black persons in the United States and an overrepresentation of Hispanic persons in the United States. The representativeness of the trial population is shown in Table S1 in the Supplementary Appendix.

END-POINT RESULTS

Figure 2.



Primary and Key Secondary End Points.

Table 2.

Table 2. Primary and Secondary End Points (Modified Intention-to-Treat Population).		
End Point	Lecanemab (N=859)	Placebo (N=875)
Primary efficacy end point		
Change from baseline to 18 mo in the CDR-SB score		
No. of participants evaluated	859	875
Adjusted mean change	1.21	1.66
Adjusted mean difference vs. placebo (95% CI)	-0.45 (-0.67 to -0.23)	
P value vs. placebo	<0.001	
Secondary efficacy end points		
Change from baseline to 18 mo in amyloid burden on PET		
No. of participants evaluated	354	344
Adjusted mean change—centiloids	-55.48	3.64
Adjusted mean difference vs. placebo (95% CI)—centiloids	-59.12 (-62.64 to -55.60)	
P value vs. placebo	<0.001	
Change from baseline to 18 mo in the ADAS-cog14 score		
No. of participants evaluated	854	872
Adjusted mean change	4.14	5.58
Adjusted mean difference vs. placebo (95% CI)	-1.44 (-2.27 to -0.61)	
P value vs. placebo	<0.001	
Change from baseline to 18 mo in the ADCOMS		
No. of participants evaluated	857	875
Adjusted mean change	0.164	0.214
Adjusted mean difference vs. placebo (95% CI)	-0.050 (-0.074 to -0.027)	
P value vs. placebo	<0.001	
Change from baseline to 18 mo in the ADCS-MCI/ADL score		
No. of participants evaluated	783	796
Adjusted mean change	-3.5	-5.5
Adjusted mean difference vs. placebo (95% CI)	2.0 (1.1 to 2.8)	
P value vs. placebo	<0.001	

Primary and Secondary End Points (Modified Intention-to-Treat Population).

The mean CDR-SB score at baseline was approximately 3.2 in both the lecanemab and placebo groups, findings consistent with early Alzheimer's disease (score of 0.5 to 6). The adjusted mean change from baseline at 18 months in the CDR-SB score was 1.21 in the lecanemab group and 1.66 in the placebo group (difference, -0.45; 95% confidence interval [CI], -0.67 to -0.23; P<0.001) (Figure 2A and Table

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



In the substudy of amyloid burden on PET (a key secondary end point) involving 698 participants, the mean amyloid level at baseline was 77.92 centiloids in the lecanemab group and 75.03 centiloids in the placebo group. The adjusted mean change from baseline at 18 months was -55.48 centiloids in the lecanemab group and 3.64 centiloids in the placebo group (difference, -59.12 centiloids; 95% CI, -62.64 to -55.60 ; $P < 0.001$) (Figure 2B and Table 2). In the modified intention-to-treat population, the mean ADAS-cog14 scores at baseline were 24.45 in the lecanemab group and 24.37 in the placebo group. The adjusted mean change from baseline at 18 months in the ADAS-cog14 score was 4.14 in the lecanemab group and 5.58 in the placebo group (difference, -1.44 ; 95% CI, -2.27 to -0.61 ; $P < 0.001$) (Figure 2C and Table 2). The mean ADCOMS in the modified intention-to-treat population at baseline was 0.398 in the lecanemab group and 0.400 in the placebo group. The adjusted mean change from baseline at 18 months in the ADCOMS was 0.164 in the lecanemab group and 0.214 in the placebo group (difference, -0.050 ; 95% CI, -0.074 to -0.027 ; $P < 0.001$) (Figure 2D and Table 2). In the modified intention-to-treat population, the mean ADCS-MCI-ADL scores at baseline were 41.2 for lecanemab and 40.9 for placebo. The adjusted mean change from baseline at 18 months in the ADCS-MCI-ADL score was -3.5 in the lecanemab group and -5.5 in the placebo group (difference, 2.0 ; 95% CI, 1.2 to 2.8 ; $P < 0.001$) (Figure 2E and Table 2).

For each of these assessments, separation of the trial groups was apparent by visual inspection of graphs at 3 months. However, no conclusions can be drawn because there was no prespecified plan for analysis that included adjustment of confidence intervals for multiple comparisons at any intermediate time point.

Sensitivity analyses of the CDR-SB score that evaluated the effect of Covid-19 (missed doses) and potential for bias from functional unblinding due to ARIA were generally consistent with the primary analysis (Table S2). Results were also consistent across key randomization strata, as well as for other factors that affect Alzheimer's disease (Figs. S1 through S4). The exploratory subgroup analysis involving ApoE $\epsilon 4$ homozygotes (15% of the trial population) numerically favored lecanemab for the ADAS-cog14 and ADCS-MCI-ADL scores but not for the CDR-SB score and the ADCOMS. Results of prespecified analyses of end points involving CSF and plasma biomarkers showed numerical improvements for all assessments comparing lecanemab with placebo, with the exception of CSF NfL (Fig. S5). In a prespecified, multiplicity-unadjusted analysis of the time to worsening of the global CDR score, the hazard ratio for progression to the next stage of dementia (0.69) numerically favored lecanemab over placebo (Fig. S6).

SAFETY

Table 3.

Table 3. Adverse Events.*

Event	Lecanemab (N=898)	Placebo (N=897)
Overall — no. (%)	798 (88.3)	735 (81.9)
Any adverse event	401 (44.7)	197 (22.0)
Adverse event related to lecanemab or placebo†	126 (14.0)	101 (11.3)
Serious adverse event	6 (0.7)	7 (0.8)
Death	62 (6.9)	26 (2.9)
Adverse event leading to discontinuation of the trial agent	237 (26.4)	66 (7.4)
Adverse event that occurred in ≥3% of participants in either group		
Infusion-related reaction	126 (14.0)	69 (7.7)
ARIA with microhemorrhages or hemosiderin deposits	113 (12.6)	15 (1.7)
Headache	100 (11.1)	73 (8.1)
Fall	93 (10.4)	86 (9.6)
Urinary tract infection	78 (8.7)	82 (9.1)
Covid-19	64 (7.1)	60 (6.7)
Back pain	60 (6.7)	52 (5.8)
Arthralgia	53 (5.9)	62 (6.9)
Superficial siderosis of central nervous system	50 (5.6)	22 (2.5)
Dizziness	49 (5.5)	46 (5.1)
Diarrhea	48 (5.3)	58 (6.5)
Anxiety	45 (5.0)	38 (4.2)
ARIA‡		
ARIA-E — no. (%)	113 (12.6)	15 (1.7)
Symptomatic ARIA-E — no. (%)§	25 (2.8)	0
ApoE ε4 noncarrier — no./total no. (%)	4/278 (1.4)	0/286
ApoE ε4 carrier — no./total no. (%)	21/620 (3.4)	0/611
ApoE ε4 heterozygote	9/479 (1.7)	0/478
ApoE ε4 homozygote	13/141 (9.2)	0/133
ARIA-E according to ApoE ε4 genotype — no./total no. (%)		
ApoE ε4 noncarrier	15/278 (5.4)	1/286 (0.3)
ApoE ε4 carrier	98/620 (15.8)	14/611 (2.3)
ApoE ε4 heterozygote	52/479 (10.9)	9/478 (1.9)
ApoE ε4 homozygote	46/141 (32.6)	5/133 (3.8)
ARIA-H — no. (%)	155 (17.3)	83 (9.3)
Microhemorrhage	126 (14.0)	68 (7.6)
Superficial siderosis	50 (5.6)	21 (2.3)
Macrohemorrhage	5 (0.6)	1 (0.1)
Symptomatic ARIA-H‡	6 (0.7)	2 (0.2)
Isolated ARIA-H: no concurrent ARIA-E	80 (8.9)	70 (7.8)
ARIA-H according to ApoE ε4 genotype — no./total no. (%)		
ApoE ε4 noncarrier	33/278 (11.9)	12/286 (4.2)
ApoE ε4 carrier	122/620 (19.7)	69/611 (11.3)
ApoE ε4 heterozygote	67/479 (14.0)	41/478 (8.6)
ApoE ε4 homozygote	55/141 (39.0)	28/133 (21.1)
ARIA-E or ARIA-H — no. (%)	191 (21.5)	85 (9.5)
Concurrent ARIA-E and ARIA-H — no. (%)	74 (8.2)	9 (1.0)

* ARIA denotes amyloid-related imaging abnormalities; ARIA-E ARIA with edema or effusions; ARIA-H ARIA with hemosiderin deposits, and Covid-19 coronavirus disease 2019.
† The relationship of adverse events to lecanemab or placebo was determined by the investigators.
‡ ARIA events were based on central review of MRI studies and include events that occurred after the double-blind intervention period.
§ Symptomatic ARIA-H concurrent with ARIA-E were included under ARIA-E.

Adverse Events.

Deaths occurred in 0.7% of the participants in the lecanemab group and 0.8% of those in the placebo group (Table 3). No deaths were considered by the investigators to be related to lecanemab or occurred with ARIA. Serious adverse events occurred in 14.0% of the participants in the lecanemab group and 11.3% of those in the placebo group. The most commonly reported serious adverse events were infusion-related reactions (in 1.2% of the participants in the lecanemab group and 0 participants in the placebo group), ARIA-E (in 0.8% and 0, respectively), atrial fibrillation (in 0.7% and 0.3%), syncope (in 0.7% and 0.1%), and angina pectoris (in 0.7% and 0). The overall incidence of adverse events was similar in the two groups (Table 3). Adverse events leading to discontinuation of the trial agent occurred in 6.9% of the participants in the lecanemab group and 2.9% of those in the placebo group. The most common adverse events (affecting >10% of the participants) in the lecanemab group were infusion-related reactions (26.4% with lecanemab and 7.4% with placebo); ARIA with cerebral microhemorrhages, cerebral macrohemorrhages, or superficial siderosis (ARIA-H; 17.3% with lecanemab and 9.0% with placebo); ARIA-E (12.6% with lecanemab and 1.7% with placebo); headache (11.1% with lecanemab and 8.1% with placebo); and falls (10.4% with lecanemab and 9.6% with placebo). Infusion-related reactions were largely mild to moderate (grade 1 or 2, 96%) and occurred with the first dose (75%). A total of 56% of the participants did not take preventative medications (i.e., nonsteroidal antiinflammatory drugs, antihistamines, or glucocorticoids) for infusion-related reactions. Of those who took preventative medications for subsequent doses, 63% did not have additional reactions.

Events of ARIA-E with lecanemab were mostly mild to moderate (91%) on the basis of central reading of imaging with the use of protocol definitions. These events were mostly asymptomatic (78%), occurred during the first 3 months of the treatment period (71%), and resolved within 4 months after detection (81%). A total of 2.8% of the participants in the lecanemab group had symptomatic ARIA-E; commonly

7.8% in the placebo group. The incidence of isolated symptomatic ARIA-H was 0.7% in the lecanemab group and 0.2% in the placebo group. The most common symptom associated with isolated symptomatic ARIA-H was dizziness. Macrohemorrhage occurred in 5 of 898 participants (0.6%) in the lecanemab group and 1 of 897 participants (0.1%) in the placebo group. ARIA-H that occurred with ARIA-E tended to occur early (within 6 months). Isolated ARIA-H occurred throughout the trial. ARIA-E and ARIA-H were numerically less common among ApoE ϵ 4 noncarriers than among carriers, with higher frequency among ApoE ϵ 4 homozygotes than among ApoE ϵ 4 heterozygotes ([Table 3](#)).

Discussion

In this phase 3 trial, the change from baseline at 18 months in the CDR-SB score (primary end point) was less with lecanemab than with placebo, favoring lecanemab. Results for secondary clinical end points were in the same direction as those for the primary end point. Lecanemab has high selectivity for soluble aggregated species of A β as compared with monomeric amyloid, with moderate selectivity for fibrillar amyloid; this profile is considered to target the most toxic pathologic amyloid species.^{4,7,8,13,14} After 18 months of treatment in the amyloid substudy, the mean amyloid level of 22.99 centiloids in the lecanemab group was below the threshold for amyloid positivity of approximately 30 centiloids, above which participants are considered to have elevated brain amyloid levels.²⁵ In the CSF substudy and in plasma analyses involving the overall population, markers of amyloid, tau, neurodegeneration, and neuroinflammation (plasma GFAP) were reduced to a greater extent with lecanemab than with placebo, with the exception of NfL, which is less sensitive to neurodegeneration than the other markers and has a slower time course for change than the others.

A definition of clinically meaningful effects in the primary end point of the CDR-SB score has not been established; however, this trial exceeded the prospectively defined target, with an estimated treatment difference of 0.373 points on a scale range of 18, a baseline value of 3.2, and early Alzheimer's disease typically characterized by a score of 0.5 to 6. In a prespecified exploratory and multiplicity-unadjusted analysis of the time to worsening (increase) of the global CDR score of at least 0.5 points on two consecutive visits, the hazard ratio for progression to the next stage of dementia numerically favored lecanemab over placebo. An open-label extension study of Clarity AD is ongoing to provide additional safety and efficacy data beyond 18 months.

In the lecanemab group, the incidence of ARIA-E was 12.6%, and the incidence of ARIA-H was 17.3%. These incidences compare with 9.9% and 10.7%, respectively, in the phase 2b trial of lecanemab, in which ApoE ϵ 4 carriers were underrepresented in the group that received 10 mg per kilogram every 2 weeks.¹⁵ The incidence of ARIA, including symptomatic ARIA, was numerically lower than in similar clinical trials, but differences in the drugs used and in trial design do not allow direct comparisons.^{26,27} ARIA-E generally occurred in the first 3 months, was mild and asymptomatic, did not lead to discontinuation of lecanemab or placebo if mild, and resolved within 4 months. The incidences of both overall and symptomatic ARIA-E were highest among ApoE ϵ 4 homozygotes.

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



Among the limitations of this trial is that it includes data for only 18 months of treatment; an open-label extension study is ongoing. The Clarity AD trial was conducted during the Covid-19 pandemic and encountered obstacles including missed doses, delayed assessments, and intercurrent illnesses. The dropout rate was 17.2%, and a sensitivity analysis that evaluated the effect of missed doses was consistent with the primary end-point analysis. An additional potential limitation was the use of modified intention-to-treat analysis without imputation of missing values. However, a sensitivity analysis that was conducted with the use of a standard intention-to-treat population with imputation yielded similar results. Finally, occurrences of ARIA may have caused participants and investigators to be aware of the trial-group assignments. We attempted to minimize this bias by making clinical raters unaware of the safety assessments and the trial-group assignments, and sensitivity analyses that were performed to examine the effect of ARIA on clinical outcomes showed that ARIA had no effect on the results. Additional trials of lecanemab include a 5-year phase 2 long-term extension trial (ClinicalTrials.gov number, [NCT01767311](#)) and a 4-year phase 3 long-term extension trial ([NCT03887455](#)) in early Alzheimer's disease, the 4-year AHEAD 3-45 trial ([NCT04468659](#)) in preclinical Alzheimer's disease, and the 4-year DIAN-TU (Dominantly Inherited Alzheimer Network Trials Unit) Next Generation trial ([NCT05269394](#)) in dominantly inherited Alzheimer's disease.

In persons with early Alzheimer's disease, lecanemab reduced brain amyloid levels and was associated with moderately less decline on clinical measures of cognition and function than placebo at 18 months but was associated with adverse events. Longer trials are warranted to determine the efficacy and safety of lecanemab in early Alzheimer's disease.

Funding and Disclosures

Supported by Eisai (regulatory sponsor), with partial funding by Biogen.

[Disclosure forms](#) provided by the authors are available with the full text of this article at NEJM.org.

This article was published on November 29, 2022, at NEJM.org.

A [data sharing statement](#) provided by the authors is available with the full text of this article at NEJM.org.

We thank the trial participants and their families, as well as all the investigators and site staff who made the trial possible (see the [Supplementary Appendix](#) for a list of collaborators); the members of the data and safety monitoring board and the raters; Lars Lannfelt and the staff of BioArctic for their early research on lecanemab; the staff of the clinical research organization Worldwide Clinical Trials for their ongoing support in conducting the trial; and J. David Cox (Mayville Medical Communications) and Lisa Yarenis (Eisai) for writing and editing assistance with an earlier version of the manuscript, in accordance with Good Publication Practice 4 ethical guidelines.

Author Affiliations

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count. 

From the Alzheimer's Disease Research Unit, Yale School of Medicine, New Haven, CT (C.H.D.); Eisai, Nutley, NJ (C.J.S., M.K., D.L., L.R., S.D., M.I., L.D.K.); the Alzheimer's Therapeutic Research Institute, University of Southern California, San Diego (P.A.); Washington University School of Medicine in St. Louis, St. Louis (R.B.); the Memory, Aging, and Cognition Center, Department of Pharmacology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore (C.C.); Eisai, Hatfield, United Kingdom (M.G.); Toronto Memory Program, Toronto (S.C.); Medical Faculty Mannheim, University of Heidelberg, Central Institute of Mental Health, Mannheim, Germany (L.F.); Katayama Medical Clinic, Okayama (S.K.), and the Department of Neuropathology, Graduate School of Medicine, University of Tokyo, and the National Center of Neurology and Psychiatry, Tokyo (T.I.) — all in Japan; Barrow Neurological Institute, Phoenix, AZ (M.S.); Toulouse Gerontopole University Hospital, Université Paul Sabatier, INSERM Unité 1295, Toulouse, France (B.V.); and Alzheimer's Research and Treatment Center, Wellington, FL (D.W.).

Dr. van Dyck can be contacted at christopher.vandyck@yale.edu or at the Alzheimer's Disease Research Unit, Division of Aging and Geriatric Psychiatry, Yale School of Medicine, 1 Church St., 8th Fl., New Haven, CT 06510.

Supplementary Material

Protocol	PDF	4873KB
Supplementary Appendix	PDF	807KB
Disclosure Forms	PDF	725KB
Data Sharing Statement	PDF	70KB

References (27)

1. Marasco RA. Current and evolving treatment strategies for the Alzheimer disease continuum. *Am J Manag Care* 2020;26:Suppl:S167-S176.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
2. Long JM, Holtzman DM. Alzheimer disease: an update on pathobiology and treatment strategies. *Cell* 2019;179:312-339.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
3. Avgerinos KI, Ferrucci L, Kapogiannis D. Effects of monoclonal antibodies against amyloid- β on clinical and biomarker outcomes and adverse event risks: a systematic review and meta-analysis of phase III RCTs in Alzheimer's disease. *Ageing Res Rev* 2021;68:101339-101339.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count. 

588.

[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

5. Sehlin D, Englund H, Simu B, et al. Large aggregates are the major soluble A β species in AD brain fractionated with density gradient ultracentrifugation. *PLoS One* 2012;7(2):e32014-e32014.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
6. Sehlin D, Hedlund M, Lord A, et al. Heavy-chain complementarity-determining regions determine conformation selectivity of anti-a β antibodies. *Neurodegener Dis* 2011;8:117-123.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
7. Magnusson K, Sehlin D, Syvänen S, et al. Specific uptake of an amyloid- β protofibril-binding antibody-tracer in A β PP transgenic mouse brain. *J Alzheimers Dis* 2013;37:29-40.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
8. Söderberg L, Johannesson M, Nygren P, et al. Lecanemab, aducanumab, and gantenerumab — binding profiles to different forms of amyloid-beta might explain efficacy and side effects in clinical trials for Alzheimer's disease. *Neurotherapeutics* 2022 October 17 (Epub ahead of print).
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
9. Hampel H, Hardy J, Blennow K, et al. The amyloid- β pathway in Alzheimer's disease. *Mol Psychiatry* 2021;26:5481-5503.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
10. Nilsberth C, Westlind-Danielsson A, Eckman CB, et al. The 'Arctic' APP mutation (E693G) causes Alzheimer's disease by enhanced A β protofibril formation. *Nat Neurosci* 2001;4:887-893.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
11. O'Nuallain B, Freir DB, Nicoll AJ, et al. Amyloid β -protein dimers rapidly form stable synaptotoxic protofibrils. *J Neurosci* 2010;30:14411-14419.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
12. Yu X, Zheng J. Polymorphic structures of Alzheimer's β -amyloid globulomers. *PLoS One* 2011;6(6):e20575-e20575.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
13. Lublin AL, Gandy S. Amyloid- β oligomers: possible roles as key neurotoxins in Alzheimer's disease. *Mt Sinai J Med* 2010;77:43-49.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



14. Haass C, Selkoe DJ. Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 2007;8:101-112.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

15. Swanson CJ, Zhang Y, Dhadda S, et al. A randomized, double-blind, phase 2b proof-of-concept clinical trial in early Alzheimer's disease with lecanemab, an anti-A β protofibril antibody. *Alzheimers Res Ther* 2021;13:80-80.
[Crossref](#) [Medline](#) [Google Scholar](#)

16. Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:270-279.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

17. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:263-269.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

18. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology* 1993;43:2412-2414.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

19. Mohs RC, Knopman D, Petersen RC, et al. Development of cognitive instruments for use in clinical trials of antidementia drugs: additions to the Alzheimer's Disease Assessment Scale that broaden its scope: the Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord* 1997;11:Suppl 2:S13-S21.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

20. Wang J, Logovinsky V, Hendrix SB, et al. ADCOMS: a composite clinical outcome for prodromal Alzheimer's disease trials. *J Neurol Neurosurg Psychiatry* 2016;87:993-999.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

21. Galasko D, Bennett D, Sano M, et al. An inventory to assess activities of daily living for clinical trials in Alzheimer's disease: the Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord* 1997;11:Suppl 2:S33-S39.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)

22. Liu KY, Schneider LS, Howard R. The need to show minimum clinically important differences in Alzheimer's disease trials. *Lancet Psychiatry* 2021;8:1013-1016

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



23. Abushakra S, Porsteinsson A, Vellas B, et al. Clinical benefits of tramiprosate in Alzheimer's disease are associated with higher number of APOE4 alleles: the "APOE4 Gene-Dose Effect". *J Prev Alzheimers Dis* 2016;3:219-228.
[Medline](#) [Google Scholar](#)
24. Insel PS, Weiner M, Mackin RS, et al. Determining clinically meaningful decline in preclinical Alzheimer disease. *Neurology* 2019;93(4):e322-e333.
[Crossref](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
25. Fleisher AS, Chen K, Liu X, et al. Using positron emission tomography and florbetapir F18 to image cortical amyloid in patients with mild cognitive impairment or dementia due to Alzheimer disease. *Arch Neurol* 2011;68:1404-1411.
[Crossref](#) [Medline](#) [Google Scholar](#)
26. Mintun MA, Lo AC, Duggan Evans C, et al. Donanemab in early Alzheimer's disease. *N Engl J Med* 2021;384:1691-1704.
[Free Full Text](#) [Web of Science](#) [Medline](#) [Google Scholar](#)
27. Budd Haeberlein S, Aisen PS, Barkhof F, et al. Two randomized phase 3 studies of aducanumab in early Alzheimer's disease. *J Prev Alzheimers Dis* 2022;9:197-210.
[Web of Science](#) [Medline](#) [Google Scholar](#)


[Close References](#)**More about**[DEMENTIA/ALZHEIMER DISEASE \(NEUROLOGY/NEUROSURGERY\)](#)[PSYCHIATRY](#)**More from the week of November 24, 2022****EDITORIAL**

ENT Drug **Universal Masking Policies in Schools and Mitigating the Inequitable Costs of Covid-19**
J. Raifman and T. Green

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.



Tap into groundbreaking research and clinically relevant insights



SUBSCRIBE

Already a subscriber? [Activate](#) your online access.

This ARTICLE IS FREE to all users and doesn't affect your subscriber-only article count.

